

Sample math exercises for students taking Physics 1600/1610 at Auburn University

Algebra

1. a) Solve the equation $x^2 - kx - 6k^2 = 0$ with respect to x , where k is a real number.
 b) Now consider the equation $x^2 - kx - 6k^2 = a$ where a is also a real number. For what values of a does this equation have no real solutions?
2. Simplify the expression

$$\frac{m_1 v_1^2}{2} + \frac{m_2 v_2^2}{2} - \frac{m_1 + m_2}{2} \left(\frac{m_1 v_1 + m_2 v_2}{m_1 + m_2} \right)^2$$

3. Solve the system of equations with respect to x and y :


$$\begin{cases} kx + ay = 2k^2 \\ 2kx + by = 3k^2 \end{cases}$$

What should the relationship between a and b be so that the system would have no solution (the equations would be incompatible)?

4. Solve the system of equations with respect to x and y :

$$\begin{cases} ax + y = \frac{1}{a} \\ xy = a \end{cases}$$

Geometry



In the setup above, a string of length l is attached to the top point A of a cylinder with the radius R . On the other end of the string there is a ball. The string touches the cylinder along the arc from point A to point B , whose angle is φ , and then goes straight, tangent to the cylinder's surface. The string and ball are on the xy -plane whose origin is at the center of the circular cross-section of the cylinder. Find the x and y coordinates of the ball in terms of R , l , and φ .

(Useful questions to ask yourself first: What length of the string touches the cylinder? What length of the string is in the air? What are the coordinates of point B ?)

Trigonometry

1. Using the half-argument identities, find the *exact* (not using a calculator, which gives you an approximate value) values of the following functions:
 $\sin 15^\circ$; $\cos 15^\circ$; $\sin 75^\circ$; $\cos 22.5^\circ$.

2. Prove the identity:

$$\frac{\cos \beta}{\cos \alpha + \sin \beta} + \frac{\sin \beta}{\cos \beta + \sin \alpha} = \frac{1 + \sin(\alpha + \beta)}{\cos(\alpha - \beta) + \frac{1}{2}(\sin 2\alpha + \sin 2\beta)}$$

3. Derive the triple-argument identities for the *sin* and *cos* functions, i. e. prove that
 $\sin 3\alpha = 3\sin \alpha - 4\sin^3 \alpha$, $\cos 3\alpha = 4\cos^3 \alpha - 3\cos \alpha$
4. Solve the equation $\sin x + \cos x = a$ for x on the interval $[0, \pi/2]$.